CLASSIFICATION RESTRICTED SSIFICATION SECURITY INFORMATION CENTRAL INTELLIGENCE AGENCY

INFORMATION FROM FOREIGN DOCUMENTS OR RADIO BROADCASTS

REPORT CD NO.

COUNTRY

German Democratic Republic

DATE OF

SUBJECT

Economic - Agriculture, cultivation zones

INFORMATION 1950 - 1951

DATE DIST.

HOW

PUBLISHED

Monthly periodicals

Oct 1952

WHERE PUBLISHED

Berlin

NO. OF PAGES 5

PUBLISHED Oct 1950, Jun 1951

SUPPLEMENT TO

REPORT NO.

LANGUAGE

German

THIS IS UNEVALUATED INFORMATION

SOURCE

Periodicals as indicated.

CULTIVATION ZONES IN THE GDR

This report consists of information from a survey of cultivation zones prepared by the GDR (German Democratic Republic) Ministry of Agriculture and Forestry as a contribution to planning for increased yields under the Five-Year Plan, and percentage data on planting of oats in the GDR, by Laender and cultivation zones.

Numbers in parentheses refer to appended sources.

Survey of Cultivation Zones

Important prerequisites for raising cultivated plants in addition to the quality of the soil and elevation, are precipitation and temperature.

To get the best yields, the following measures must be carried out:

- 1. All types of plants which are not sufficiently productive must be eliminated.
 - 2. The entire arable land must be divided into cultivation zones.
- 3. Within each cultivation zone a type-selection plan must be put into effect which uses the hereditary factors of the plants most effectively.

Therefore, the Ministry of Agriculture and Forestry worked out a weather map for the 126 Kreise of the GUR. The individual cultivation zones were determined on the basis of the following natural conditions:

- 1. Average precipitation (for about 40 years)
- 2. Yearly average and July temperatures (for about 40 years)
- 3. Elevation above sea level

-1-

CLASSIFICATION				RESTRI	CEED				
STATE X NAVY	$\exists x$	NSRB		DISTRI	BUTION				L
ARMY AIR		FBI							

STAT



RESTRICTED

- 4. Soil quality
- 5. The ratios of the various plant types sown to one another.

In order eventually to establish the cultivation zones correctly, the chemical composition of the harvested plants will be studied to determine whether the hereditary factors of individual plants have been preserved. On the basis of these results, phenological maps for smaller regions will be made.(1)

See table and map on following pages.



- 2 -

RESTRICTED


Sanitized Copy Approved for Release 2011/08/11 : CIA-RDP80-00809A000700090019-0

Characteristics of GDR Cultivation Zones (1)

50		Cultivation Zone	Region	Elev Above Sea Level (meters)	Av Precipitation (mm)	Av Temp Annual	(°C) July	<u>Description</u>	
- 3 - RKSTRICTED		I	Coast	0-100	500-650				
					J00-030	8	17.1	Poor to mediocre soil; cool summer,	
	'	II	Lowland	0-100	500- <i>6</i> 00	8.5	17.9	mild winter	
		III				•	-1.5	Poor to good soil, marshy and hurms (high ground-water level in some places); warm summer, mild winter	
			Flatland	50-200	450-550	7.8	18.3	Mostly poor soil, partly flinty, partly washed-out (low ground-water	
	IV	Fertile plains					level); warm summer, mild winter		
		pasture land	100-200	450-550	8	17.9	Very good soil (good clay and loose		
	v	Transition					soil); warm summer, mild winter		
			region	150-400	500-600	7-7	17.7	Mostly good soil; arm summer, cold	
	AI	Foothills	200-450	500-600	7.4	17.5	winter Mediocre soil; warm summer, cold		
	VII	Mountains	450-1,000	4-1		_,.,	winter		
				+70-1,000	650-1,000	5.8	16.5	Agriculturally poor soil (some better soil in valleys); cool sum- mer, severe winter	

STAT

Sanitized Copy Approved for Release 2011/08/11: CIA-RDP80-00809A000700090019-0



RESTRICTED

STAT



RESTRICTED

Planting of Oats in the GDR

Γ

In the GIR, 10.7 percent of the arable land is used for raising oats. The following percentages of arable land in the Laender are used for raising oats: Brandanburg 9.3, Sachsen-Anhalt 9.3, Thueringen 11, Macklenburg 11.3, and Sachsen 14:

Of the total cats acreage in the GDR, Mecklenburg has 23.9 percent, Sachsen-Anhelt 23.1, Sachsen 20.6, Brandenburg 17.8, and Thueringen 14.6.

The following table shows the percentage distribution of various types of cats among the Laender:

Types of Oats	Branden- burg	Mecklen- burg	Sachsen- Anhalt	rm .	_
Flaemingsgold			Author 6	Thueringen	Sachsen
_	22	28	15	13	22
Flaemingstreue	26	7	28	2	37
PSG Goldkorn	31	9	22	29	9
Svalcefs Goldregen III	3	3	8	7	79
Kirsches Pfiffelbacher		1	10	·	,,,
		•	19	59	21
Dippes Weiss	9	10	35	- 5	41
Siegeshafer	27	45	20		8

If the area of the GDR is broken down according to cultivation zones, the following percentages of each zone's arable land are used for raising oats:

Cultivation Zone	Arable Land Used for Oat Raising (%)
I Coast	11.7
II Lowland	10.7
III Flatland	9.2
IV Fertile plains	8.5 ·
V Transition region	12.5
V1 Foothills	14.0
VII Mountains	15.2
All GDR	10.7 (2)

SOURCES

- 1. Berlin, Die Deutsche Landwirtschaft, Oct 50
- 2. Berlin, Statistische Praxis, Jun 51

- E N D -

-5-

RESTRICTED

STAT

